What Happens When You Consume a Cup of Coffee?

By Joachim Bartoll | Jan. 26th, 2026

Send to Kindle

Ah, the modern indoctrinated slave's siren call: that first sip of coffee, the elixir of "getting through the day." Picture the average zombie shuffling to the kitchen, eyes half-open, typing the typical message on social media, "Can't go to work without my coffee jolt, need that caffeine kick to fire up!" Little do they know, they're not "waking up," they're igniting a full-body panic attack, courtesy of plant poisons masquerading as productivity. That dragging fatigue deceiving them into this harmful backwards habit? Not from life's cruelties, but from yesterday's cups, a crappy plant-laden diet, and a lifestyle that screams for real nourishment. Meanwhile, their body begs for mercy, but hey, who has time for that when there's another stress spike brewing? Let's dive into the science of this daily disaster, where "perk up" really means "poison up."

As you should know by now, in the realm of human nutrition and physiology, grounded in anthropology, biology, and biochemistry, it is undeniable evident that our species has evolved as hypercarnivores, thriving on animal-based foods that provide bioavailable organic nutrients in chemical organic forms identical to those stored in our own bodies. Plant-derived inorganic substances, however, are not only chemically incompatible, they often carry a burden of defense chemicals, antinutrients, heavy metals, and pesticide residues, which are extremely toxic to our biological terrain. These elements, designed by plants to deter predators, introduce toxicity and disrupt our internal environment, leading to a compromised terrain that manifests as modern ailments. Coffee, derived from the seeds of the Coffea plant (the most toxic part of any plant,) exemplifies this mismatch. While it may offer a fleeting sense of alertness from a stress

response, its consumption also unleashes a cascade of physiological disruptions that burden the body, far outweighing any perceived benefits. This article delves into the precise, step-by-step processes occurring in the human body upon ingesting 177 milliliters of typical brewed coffee, equivalent to a standard six-ounce cup. I will draw from established scientific literature in endocrinology, human physiology, biochemistry, and microbiology, emphasizing the terrain theory, which posits that disease arises not from external pathogens but from an imbalanced internal milieu exacerbated by toxins and stimulants. By understanding these mechanisms, we can appreciate the wisdom of aligning with our species-appropriate diet of raw or lightly cooked animal foods, free from such interferences, to foster true vitality and longevity.

The journey of coffee through the body commences the instant it contacts your lips, marking the onset of a series of physiological interactions that reveal its inherent incompatibility with human biology. As a typically hot and inherently bitter liquid, coffee first engages with the sensitive oral mucosa and specialized taste receptors on the tongue. This bitterness, primarily derived from diterpenes such as cafestol and kahweol, functions as an biological warning mechanism signaling potential harm to any organism attempting to consume it. This is why most normal people hate the taste of coffee the first time they try it, as their body instinctively warns them and wants you to throw it away, but then people are conditioned through social norms to drink it anyway, to suppress their instincts and harm their bodies. Insanity. Now, these diterpenes, alongside alkaloids including caffeine, theobromine, and theophylline, represent potent toxins synthesized by the Coffea plant to deter insects, herbivores, and other threats in its natural environment. They are biochemical weapons developed by the plant to maim, sterilize and kill. Caffeine, in particular, evolved as a natural insecticide, leaching into surrounding soil to inhibit competing organisms and protect the seeds, underscoring its role not as a benign stimulant but as a biochemical weapon. In humans, this initial encounter elicits minor yet telling sensory

responses, such as heightened salivation, as the body instinctively seeks to dilute and neutralize the incoming irritants through increased fluid secretion.

However, beneath this superficial reaction lies a more insidious concern: the subtle yet cumulative damage inflicted upon the delicate epithelial lining of the mouth and esophagus. The combination of coffee's elevated temperature and its acidic nature, characterized by a pH typically around 5, can induce micro-abrasions, localized inflammation, and even cellular destruction in these tissues. Heat alone has the potential to compromise the integrity of the esophageal mucosa, necessitating repeated cellular regeneration that, over time, may predispose the area to chronic vulnerabilities. Compounding this, antinutrients like tannins and chlorogenic acids initiate their disruptive actions here, binding avidly to proteins in saliva and potentially diminishing enzymatic activity right from this early stage. Tannins, in particular, have been shown to significantly impair the lubricating properties of saliva, leading to a drier oral environment that exacerbates irritation and reduces the natural protective barrier against toxins. Chlorogenic acids, major polyphenolic compounds in coffee, further interact with salivary cells and proteins, contributing to astringency and potential discoloration of dental enamel through their binding affinity, while also promoting oxidative stress in the oral cavity. Caffeine itself may reduce overall salivary flow, fostering xerostomia or dry mouth, which impairs the mouth's ability to buffer acids and clear debris, thus amplifying the toxic burden. This is especially bad among people who consume anything plant based, such as carbohydrates, that stick to the teeth and attract the wrong kind of bacteria that damages the enamel. With a reduction in salivary flow from the toxins in coffee, the damage on the teeth is multiplied if you eat something close to or after having a cup of coffee, as most people do.

As the liquid progresses downward through the esophagus, propelled by coordinated peristaltic waves toward the stomach, the embedded stimulants begin signaling the autonomic nervous system, triggering a preliminary

sympathetic activation that foreshadows the broader stress response to come. This early neural alert diverts physiological resources toward defense/detoxification, even before full absorption, highlighting coffee's role as an exogenous stressor misaligned with our human physiology. This entire phase unfolds in mere seconds, yet it signifies the body's primal recognition of coffee as a foreign invader, laden with elements antithetical to our biological terrain.

From the perspective of terrain theory and human physiology, these plant-derived chemicals introduce a host of disruptive agents that erode internal homeostasis. Notably, coffee carries deuterium-rich components, as plant foods generally exhibit higher deuterium levels compared to animal-based sustenance, with concentrations elevated in seeds like coffee beans due to their metabolic storage patterns. This heavy hydrogen isotope interferes with the cell's mitochondrial efficiency, subtly impairing cellular energy production and fostering an acidic milieu conducive to imbalance rather than equilibrium. Compounding this, heavy metals such as lead and cadmium frequently contaminate coffee beans through soil uptake, agricultural practices, and processing, with studies detecting these toxins in virtually all tested samples. Lead and cadmium accumulate in the body, disrupting enzymatic functions and mitochondrial integrity, while also contributing to systemic toxicity that aligns with the compromised terrain underlying modern degenerative conditions. Additionally, residual pesticides from non-organic cultivation, including synthetic herbicides and fungicides, add to this load, further taxing the oral and esophageal tissues upon contact and setting the stage for broader detoxification demands. In anthropology and biology, our pre-agricultural ancestors, adapted to a diet scarce in edible plants and rich in bioavailable animal nutrients, possessed no evolutionary preparation for such assaults, rendering these interactions a profound mismatch that initiates a cascade of physiological strain from the very first sip.

References: Esophageal Cancer Risk: The Truth About Hot Drinks Like Coffee and Tea, The surprising connection between your morning coffee and the battle between plants and herbivory insects – Boyce Thompson Institute, Saliva tannin interactions – PubMed, Influence of Coffee on Soft & Hard Dental Tissues – MyZerodonto, Evaluation of chlorogenic acids and melanoidin interactions with salivary proteins and their effect on tooth discoloration – ScienceDirect, Deuterium Content in different foods and Deuterium Depletion – Osmio Water Filters, Deuterium in Histamine Intolerance, Coffee Study – Clean Label Project, Heavy Metals in Coffee: What You Need To Know – Koffee Kult, Toxins in Coffee: Should You Be Worried? Here's What to Know – Pure Essentials Supplements

Following the transition of coffee's compounds into the stomach, they confront the gastric environment tailored for processing bioavailable nutrients from animal sources. The human stomach is solely constructed to efficiently break down animal proteins and fats using hydrochloric acid and enzymes such as pepsin, reflecting our anthropological heritage as hypercarnivores adapted to a diet scarce in edible plants before the detrimental agricultural revolution. This physiological setup renders the stomach ill-equipped to handle the influx of plant-derived toxins and antinutrients inherent in coffee, which disrupt its natural functions and introduce incompatibility with our species-appropriate biology. Animal proteins, being complete and highly digestible with superior amino acid bioavailability compared to plant sources, align seamlessly with gastric mechanisms, whereas plant toxins like those in coffee provoke defensive responses that strain the system. Coffee's inherent acidity, with a pH typically around 5, exacerbates this mismatch by stimulating gastrin release from G cells in the stomach and duodenum, thereby increasing hydrochloric acid secretion by up to 30 percent. Caffeine plays a central role here, activating bitter taste receptors on parietal cells to further induce gastric acid secretion through

complex signaling pathways, independent of its stimulant effects elsewhere in the body. Both regular and decaffeinated coffee prompt a prompt and sustained elevation in serum gastrin levels, confirming that non-caffeine components also contribute to this hypersecretory response. This surge in acidity can lead to hyperacidity, progressively eroding the protective mucus layer that lines the stomach walls and safeguards against autodigestion, thus exposing the underlying epithelial cells to direct irritation and potential ulceration. The mucus barrier, composed of mucins secreted by goblet cells, is crucial for maintaining gastric homeostasis, but coffee's compounds weaken this defense, increasing vulnerability to damage. Compounds like chlorogenic acids and caffeoylquinic acids, which are polyphenolic defense chemicals abundant in coffee beans, further aggravate this process by inhibiting mucin production and promoting local inflammation through oxidative stress and interaction with mucosal cells. These acids, even in decaffeinated varieties, have been linked to increased stomach discomfort, including dyspepsia, nausea, and heartburn, by elevating gastric acid levels and irritating the lining, underscoring their role as irritants rather than benign elements.

Enzymatic activity within the stomach also suffers significantly under this assault. The high amounts of tannins (antinutrients) in coffee beans bind tightly to digestive enzymes and proteins, reducing their efficacy and slowing the breakdown of ingested materials. These condensed polyphenols inhibit key enzymes such as trypsin, amylase, and lipase by altering their substrate binding affinity, a mechanism that extends beyond the stomach to impair overall digestion. Phytic acid, another prominent antinutrient in coffee, chelates essential minerals like zinc and iron, forming insoluble complexes that prevent their absorption and utilization while causing tissue build-ups, exacerbating deficiencies already prevalent in diets incorporating plant foods where such minerals are inorganic and much less bioavailable. This chelating property stems from phytic acid's phosphate groups, which avidly bind divalent cations,

inhibiting not only mineral uptake but also enzymes like pepsin required for protein digestion. Oxalates, present in coffee, form insoluble crystals that can irritate the gastric mucosa, contributing to pain, inflammation, and potential disruption of the gut barrier. These crystals, by binding calcium and other minerals, will heighten oxidative stress and mucosal damage. They also form kidney stones and compromise connective tissues. Melanoidins, generated during the roasting process as end products of the Maillard reaction between amino acids and reducing sugars, add to the toxic load with their brown, nitrogen-containing, high-molecular-weight structures that exhibit carcinogenic properties through promotion of oxidative stress and severe damage. Pesticide residues, including glyphosate and pyrethroids from conventional farming practices, linger in coffee beans and amplify this damage by disrupting the gut microbiota balance, as evidenced by microbiology studies showing alterations in microbial composition and reduced beneficial bacteria. Glyphosate, in particular, targets the shikimate pathway in gut microbes, leading to dysbiosis that impairs digestion, increases permeability, and heightens systemic toxicity. The immediate outcomes of these interactions often manifest as discomfort, such as bloating or nausea, stemming from gas production and impaired motility, but more insidiously, they compromise the stomach's barrier function, causing leaky gut and permitting undigested particles and toxins to leak into the bloodstream and propagate further physiological strain.

In terms of terrain theory, this phase markedly acidifies the biological terrain, creating an internal milieu where cellular debris accumulates, pH imbalances prevail, and detoxification pathways become overburdened, deviating from the balanced environment fostered by animal-based nutrition. For a hypercarnivore, whose stomach thrives on the neutral pH and nutrient-dense profile of animal fats and proteins, this intrusion represents a profound mismatch, leading to cumulative erosion of mucosal integrity and overall vitality over time.

References: A Metabolomic Perspective on Carnivorous and Scavenger Adaptations – MDPI, Alternative proteins vs animal proteins: The influence of structure and processing on their gastro-small intestinal digestion – Science Direct, Caffeine induces gastric acid secretion via bitter taste signaling in gastric parietal cells – PubMed Central, Effect of regular and decaffeinated coffee on serum gastrin levels – PubMed, Effects of Coffee on the Gastro-Intestinal Tract – MDPI, The acute effects of coffee and caffeine on human interdigestive exocrine pancreatic secretion – Mayo Clinic, Caffeine induces gastric acid secretion via bitter taste signaling – PNAS, Gastric acid secretion, gastrin release, and gastric emptying in humans as affected by liquid meal temperature – ScienceDirect, Effect of Regular and Decaffeinated Coffee on Serum Gastrin Levels – Gwern.net, Effects of Coffee on the Gastro-Intestinal Tract – PubMed, Why Can Coffee Cause Stomach Pain? – Vinmec, An immobilized fungal chlorogenase rapidly degrades chlorogenic acid in a coffee beverage without altering its sensory properties – ScienceDirect, Chlorogenic Acid–Gut Microbiota Interactions: New Insights for Alleviating Chronic Diseases and Health Disorders – Oxford Academic, Inhibition of in vitro enzymatic starch digestion by coffee extract – ScienceDirect.com, Inhibitory effect and mechanism of tannic acid against two starch digestive enzymes – Oxford Academic, Tannins From Different Foodstuffs As Trypsin Inhibitors – Institute of Technical Biochemistry, The inhibition of digestive enzymes by polyphenolic compounds – PubMed, Slowing starch digestion and inhibiting digestive enzyme activity using plant flavanols/tannins – ScienceDirect, The effect of concentration of tannin-rich bean hulls on activities of lipase and α-amylase – Cambridge, Inhibition of Trypsin by Condensed Tannins – Journal of Agricultural and Food Chemistry, Inhibitory effect and mechanism of tannic acid against two starch digestive enzymes – Oxford Academic, Reduction of phytic acid and enhancement of

bioavailable micronutrients in food grains, Characterization of a Metal-Chelating Substance in Coffee – ResearchGate, Minerals (Zn, Fe, Ca and Mg) and Antinutrient (Phytic Acid) Constituents in Common Bean – PMC – NIH, Living With Phytic Acid – The Weston A. Price Foundation, The Dangers of Phytic Acid and What to Do About It | Embracing Motherhood, Phytic Acid: An Optimal Barrier to Iron Absorption, Know Your Poison: Phytic Acid – Joachim Bartoll, Coffee Consumption With The Incident Of Gastritis – Journal of Applied Nursing and Health, Oxalates: The IBS Trigger No One is Talking About – Amaluna Wellness, Why High Oxalates Can Harm Your Health | MosaicDX – Mosaic Diagnostics, Oxalates Fact Sheet | High Oxalate Foods – Melbourne Functional Medicine, Know Your poison: Oxalates – Joachim Bartoll, Assessment of Healthy and Harmful Maillard Reaction Products in a Novel Coffee Cascara Beverage – PubMed Central, Exploring Maillard reaction melanoidins to investigate toxicological profiles of snacks from corn/common bean mixtures, Effects of Maillard reaction and its product AGEs on aging and age-related diseases, Chemical Composition and Potential Biological Activity of Melanoidins From Instant Soluble Coffee and Instant Soluble Barley – NIH, Is Coffee Full of Pesticides? What NZ Drinkers Should Know, Pesticides: Unintended Impact on the Hidden World of Gut Microbiota, How Pesticides in Your Brew May Harm Your Gut Health, A review on the pesticides in coffee – NIH, Pyrethroid and metal residues in different coffee beans, Effects of Coffee on the Gastro-Intestinal Tract – NIH, Coffee, Acid Reflux, and GERD, Kinetic characterization of thermophilic and mesophilic anaerobic digestion for coffee grounds and waste activated sludge – ScienceDirect, Acids in coffee: A review of sensory measurements and meta-analysis of chemical composition – Taylor & Francis, A Meta-Analysis of Acids in Coffee and a Quantification of Coffee Beverage

> Color – ProQuest, Relationship between the Chemical Composition and the Biological Functions of Coffee

Following its passage through the stomach, the partially digested coffee chyme reaches the small intestine by approximately 30 to 45 minutes post-ingestion, entering an environment optimized for the absorption of bioavailable nutrients from animal-based foods, as per our hypercarnivorous physiological adaptations rooted in anthropology and biology. This phase marks a significant intensification of absorption, where the small intestine's extensive surface area, provided by villi and microvilli, facilitates the rapid uptake of coffee's various compounds into the bloodstream. However, unlike the organic, bioactive forms of nutrients stored in animal tissues that align with human biochemistry, coffee's plant-derived substances introduce an array of stimulants, defense chemicals, antinutrients, heavy metals, and pesticide residues that disrupt this process, diverting energy toward neutralization and detoxification. The duodenum, jejunum, and ileum sequentially handle this influx, with most absorption occurring in the jejunum due to its high density of transporters and enzymes, though our evolutionary lack of specialized machinery for plant toxins exacerbates the burden.

Caffeine, the predominant stimulant in coffee at roughly 100 milligrams per standard cup, undergoes swift absorption primarily through the intestinal villi, achieving nearly 99 percent uptake within 45 minutes via passive diffusion across the lipid membranes of enterocytes. This rapid process is facilitated by caffeine's lipophilic nature, allowing it to cross the brush border without requiring active transport, though it can also utilize organic cation transporters in cases of lower concentrations. Once absorbed, caffeine enters the portal vein and circulates systemically, but its entry triggers immediate disruptions. Theobromine and theophylline, fellow methylxanthines present in lesser amounts, follow a comparable pathway, crossing into the bloodstream via passive diffusion and

reaching peak plasma levels within one to two hours. These compounds, structurally similar to caffeine, are absorbed at rates of 90 to 100 percent, with theobromine exhibiting a slightly slower uptake due to its lower solubility, yet still contributing to the overall stimulant load. Collectively, these methylxanthines antagonize adenosine receptors in the central nervous system and periphery, inhibiting natural inhibitory signals that promote relaxation and instead enforcing a stressful state of heightened alertness. This blockade incurs a substantial physiological cost, as it prompts the activation of the hypothalamic-pituitary-adrenal axis, leading to the release of stress hormones such as cortisol and adrenaline, thereby initiating the fight-or-flight response that diverts resources from restorative processes. In the context of endocrinology, this artificial stimulation mimics a threat, elevating heart rate, blood pressure, and glucose mobilization, while suppressing non-essential functions like digestion, detoxification and tissue repair. In other words, your body readies itself for fight or flight and all tissue repair and healing is shut down, accelerating any kind of toxic build-up, tissue damage and thus aging.

Defense chemicals such as the diterpenes cafestol and kahweol, inherent to coffee beans as protective agents against herbivores, are absorbed less efficiently but nonetheless enter circulation, with studies indicating 70 to 90 percent uptake in the small intestine, primarily in the ileum where lipid absorption predominates. These lipophilic compounds, often esterified with fatty acids in unfiltered coffee, require micellar solubilization by bile salts for emulsification and subsequent incorporation into chylomicrons for lymphatic transport, bypassing initial hepatic metabolism and allowing direct systemic distribution. Due to poor hepatic metabolism, involving limited cytochrome P450 activity, cafestol and kahweol accumulate in tissues, interfering with cholesterol homeostasis by inhibiting enzymes like cholesterol ester transfer protein and elevating LDL levels as the body attempts compensatory repair. Their persistence

exacerbates oxidative stress and tissue damage, misaligning with human physiology adapted to minimal plant lipid toxins.

Antinutrients like tannins and oxalates further compound the disruption by binding to intestinal transporters and impairing the absorption of essential minerals and proteins. Tannins, polyphenolic compounds, form complexes with divalent cations such as iron and zinc, reducing their bioavailability by up to 50 percent in the jejunum, while also inhibiting brush border enzymes like sucrase and maltase, slowing carbohydrate breakdown and contributing to osmotic imbalances. Slowing down carbohydrate breakdown and glucose release is extremely bad as that results in prolonged elevated glucose levels in the blood with increased glycation, the binding of glucose to fatty acids and protein, rendering them very harmful and contributing to cardiovascular- and heart disease.
Oxalates, meanwhile, chelate calcium and magnesium, forming insoluble crystals that not only hinder mineral uptake but also induce oxidative stress on enterocytes through reactive oxygen species generation, damaging mitochondrial function and compromising the epithelial barrier. This stress manifests as lipid peroxidation and reduced glutathione levels, fostering an inflammatory milieu that weakens tight junctions and increases permeability, allowing further toxin leakage.

Heavy metals, including lead, cadmium, and arsenic, concentrated in coffee beans due to soil contamination and agricultural practices, are partially absorbed here, with cadmium exhibiting high bioavailability via zinc transporters in the duodenum, leading to accumulation in kidneys and liver where it induces metallothionein binding but overwhelms detoxification at chronic levels. Lead competes with calcium for absorption, disrupting neuronal signaling and contributing to systemic toxicity, while arsenic, often from groundwater irrigation, methylates inefficiently in humans, promoting oxidative damage and carcinogenesis. Pesticide residues, such as glyphosate and pyrethroids, lingering

in the bean's seed, undergo partial absorption through passive diffusion and active transport, with glyphosate inhibiting the shikimate pathway in gut microbes and pyrethroids altering sodium channels, both amplifying oxidative stress and endocrine disruption. Microbiology insights reveal that these compounds profoundly alter the gut microbiome, selectively reducing beneficial bacteria like Bifidobacterium and Lactobacillus while promoting pathogenic overgrowth of fiber and starch fermenting bacteria, leading to dysbiosis that impairs short-chain fatty acid production, weakens the mucosal barrier, causes barrier tissue damage, further compromising the biological terrain. In line with terrain theory, this dysbiotic shift acidifies the intestinal environment, accumulating cellular debris and overburdening systemic detoxification.

This absorption phase culminates around one hour, inundating the body with incompatible substances that our pre-agricultural ancestors, subsisting on a diet devoid of such concentrated plant toxins, never encountered. Anthropology indicates that humans, adapted to scarce edible plants amid famine, lacked the enzymatic machinery, such as enhanced cytochrome P450 variants or specialized gut flora, to efficiently handle these loads, as evidenced by cell studies showing limited to non-existent adaptations to post-agricultural plant defenses. Consequently, energy is immediately redirected toward detoxification pathways, including phase I oxidation and phase II conjugation, rather than toward nourishment and repair, underscoring the profound mismatch with our species-appropriate animal-based diet.

References: Effects of caffeine and coffee on fluid transport in the small intestine – Gastroenterology, Emptying and Absorption of Caffeine from the Human Stomach – ScienceDirect, The Competitive Absorption by the Gut Microbiome Suggests the First-Order Absorption Kinetics of Caffeine – Sage Journals, Theobromine, caffeine, and theophylline metabolites in human plasma and urine – ScienceDirect, Psychopharmacology of theobromine in healthy volunteers – PMC – PubMed Central,

Health Effects and Risks of Caffeine, Theobromine and Theophylline | Bentham Science, Theobromine | C7H8N4O2 | CID 5429 – PubChem – NIH, Bioavailability and Potential Impact on Human Health of Caffeine, Theobromine, and Trigonelline – Wiley Online Library, Effect of Coffee Lipids (Cafestol and Kahweol) on Regulation of Cholesterol Metabolism, Separate effects of the coffee diterpenes cafestol and kahweol on serum lipids and liver aminotransferases – PubMed, Cafestol and Kahweol: A Review on Their Bioactivities and Pharmacological Properties, On the Cholesterol Raising Effect of Coffee Diterpenes Cafestol and 16-O-Methylcafestol – MDPI, Tannic acid attenuates intestinal oxidative damage – PMC – NIH, Analysis of toxic heavy metals in coffee samples – ACS, Heavy metal and mycotoxin-producing fungi contamination of coffee consumed in Saudi Arabia – ScienceDirect, Assessment of the risk of exposure to cadmium and lead as a result of the consumption of coffee infusions – PMC – NIH, Heavy-Metal Contents and the Impact of Roasting on Polyphenols, Caffeine, and Acrylamide in Specialty Coffee Beans – MDPI, A quantitative assessment of heavy metal contamination in instant coffee beverages – Polish Platform of Medical Research, A quantitative assessment of heavy metal contamination in instant coffee beverages, A review on the pesticides in coffee: Usage, health effects, detection, and mitigation – NIH, Study of Pesticide Mixtures and Intestinal Barrier Dysfunction Further Highlights Regulatory Deficiencies, The Coffee Paradox: How Pesticides in Your Brew May Harm Your Gut Heal, Pyrethroid and metal residues in different coffee bean preparing processes and their human health risk assessments via consumption | bioRxiv, Effects of Coffee and Its Components on the Gastrointestinal Tract and the Brain–Gut Axis – PMC – NIH, Coffee Consumption Modulates Amoxicillin-Induced Dysbiosis – Frontiers, The Association between Caffeine Intake and the Colonic Mucosa – MDPI, Toxic substances in plants and the

> food habits of early man – PubMed, Noah Whiteman digs into the plant toxins we consume – C&EN, Noah Whiteman: Exploring the Chemistry of Plant Toxins and Human Culture, Plant Toxins, Accumulated Toxicity, Tissue Damage, And a Decreased Lifespan – Joachim Bartoll, Antinutrients? Everything In Vegetables Is Unusable And Toxic To Humans! – Joachim Bartoll, How Plants Make, Store and Use Toxins – Learn Genetics Utah, Plant Toxic Proteins: Their Biological Activities, Mechanism of Action – ResearchGate

Upon absorption in the small intestine, coffee's myriad compounds, including stimulants, defense chemicals, antinutrients, heavy metals, and pesticide residues, enter the bloodstream and commence their systemic circulation, rapidly disseminating throughout the body to exert widespread physiological disruptions incompatible with our human biology and physiology. This phase begins immediately after uptake, with compounds reaching peak plasma levels within 45 to 60 minutes, a timeframe that underscores the swift infiltration of these plant-derived toxins into vital tissues and organs. Caffeine, the primary alkaloid at approximately 100 milligrams per cup, exhibits a distribution volume of about 0.7 liters per kilogram of body weight, indicating its hydrophilic nature and free diffusion into intracellular water across all tissues, including those ill-equipped to handle such foreign invaders. Its analogs, theobromine and theophylline, follow suit, permeating the bloodstream and achieving widespread distribution due to their lipophilic properties, allowing passive diffusion across cellular membranes. Defense chemicals like the diterpenes cafestol and kahweol, though less efficiently absorbed, still enter circulation via lymphatic transport in chylomicrons, bypassing initial hepatic clearance and facilitating direct systemic spread, where their poor metabolism leads to prolonged presence and accumulation in tissues. Antinutrients such as tannins and oxalates, along with toxins like melanoidins and acrylamide formed during roasting, circulate bound to plasma proteins or in free form, promoting oxidative stress and tissue damage

systemically. Heavy metals including lead, cadmium, and arsenic, concentrated in coffee beans, distribute via blood to accumulate in organs like the liver and kidneys, exacerbating toxicity. Pesticide residues, such as glyphosate, similarly spread, disrupting endocrine signaling and cellular function across the body.

Caffeine and its analogs readily cross the blood-brain barrier due to their liposolubility, stimulating the central nervous system by antagonizing adenosine receptors and thereby preventing natural inhibitory signals that promote relaxation. This central action activates the hypothalamic-pituitary-adrenal (HPA) axis, a key stress regulatory pathway, through interactions with centrally located adenosine receptors in hypothalamic afferent regions, eliciting a full stress response akin to a perceived threat. The adrenal glands consequently release cortisol, with levels increasing by 25 to 100 percent, and adrenaline, mimicking an external danger and preparing the body for fight-or-flight. This hormonal surge accelerates heart rate by stimulating beta-adrenergic receptors, raises blood pressure through vasoconstriction and increased cardiac output, and mobilizes glucose from hepatic stores via glycogenolysis and gluconeogenesis to provide immediate energy. However, this artificial surge, devoid of a genuine threat, depletes energy reserves, leading to rebound fatigue as the body compensates for the unwarranted mobilization, fostering a cycle of dependency and exhaustion.

In the context of terrain theory, this disrupts the biological terrain by introducing an acidic, deuterium-laden milieu from plant sources, overburdening cellular energy production and favoring imbalance.

Endocrinology further illuminates the hormonal disruptions precipitated by this systemic spread. Elevated cortisol, a glucocorticoid, suppresses thyroid function by inhibiting the conversion of thyroxine (T4) to active triiodothyronine (T3,) reducing free T3 levels and impairing metabolic rate, which aligns with observations of altered thyroid hormone profiles in caffeine consumers. Chronic

stimulation from repeated exposure fosters insulin resistance, as cortisol antagonizes insulin action, elevating blood glucose and promoting fat storage, particularly in a terrain compromised by plant toxins and high glucose from non-animal sources. Diterpenes like cafestol and kahweol interfere with lipid metabolism by upregulating cholesteryl ester transfer protein (CETP) and phospholipid transfer protein (PLTP,) disrupting the heteroexchange of lipids between lipoproteins such as VLDL, HDL, and LDL, thereby elevating LDL cholesterol as the body ramps up production for repair of toxin-induced damage. In a toxic terrain, exacerbated by deuterium-rich coffee and pesticide residues, this elevation fosters oxidized LDL particles and inflammation, as damaged lipids accumulate without efficient recycling. Toxins such as melanoidins, high-molecular-weight Maillard reaction products from roasting, and acrylamide, a neurotoxin and carcinogen, promote oxidative stress by generating reactive oxygen species, damaging endothelial cells through lipid peroxidation and impairing nitric oxide bioavailability, thus contributing to cardiovascular strain including atherosclerosis and hypertension. Acrylamide, in particular, induces endothelial dysfunction and increases cardiovascular risk via inflammation and transforming growth factor-beta1 pathways, while melanoidins, despite some purported antioxidant effects in isolation, add to the overall toxic load in the context of coffee's chemical cocktail.

From a physiological standpoint, this distribution imposes a substantial burden on key organs, particularly the heart and brain, where the stimulants and toxins exert direct and insidious effects. In the cardiovascular system and the heart, caffeine and adrenaline cause vasoconstriction of coronary arteries, temporarily elevating blood pressure and heart rate, which strains cardiac muscle and increases myocardial oxygen demand, potentially exacerbating arrhythmias or ischemia in a compromised terrain. Accumulated diterpenes and heavy metals further damage cardiovascular tissue, promoting fibrosis and reduced elasticity. In the brain, neuronal overexcitation from adenosine blockade leads to

heightened alertness but also risks excitotoxicity, contributing to anxiety through amplified sympathetic activity and cortisol feedback, as well as headaches from cerebral vasoconstriction and rebound vasodilation upon withdrawal. This overexcitation mimics chronic stress, depleting neurotransmitters and fostering neuroinflammation, inconsistent with our evolutionary adaptation to a low-toxin, animal-based diet that supports stable neural function without such artificial perturbations.

References: Pharmacology of caffeine and its effects on the human body – ScienceDirect, The Effects of Caffeine on Your Body | Blog – ProMedica, Caffeine as a Factor Influencing the Functioning of the Human Body – PMC, The Effects of Caffeine on Your Body – ProMedica,Caffeine as a Factor Influencing the Functioning of the Human Body – PMC,Caffeine is a Defense Chemical and a Potent Poison — Do Not Neglect Accumulated Damage – Joachim Bartoll,The Accumulated Damage – Daily Caffeine Use – Joachim Bartoll,Caffeine and Peripheral Blood Flow – Auctores,Caffeine metabolism and heart health – FitnessGenes,Coffee Acutely Increases Sympathetic Nerve Activity and Blood – AHA Journals,Hypothalamo-pituitary-adrenocortical axis – Caffeine modulation – PMC,Exploring the Impact and Mechanisms of Coffee and Its Active – MDPI,Neurocognitive and Neurological Effects of Coffee and Caffeine – Cureus,Caffeine Consumption – Depression, Anxiety, and Stress Review – Springer,Caffeine Stimulation of Cortisol Secretion Across the Waking Hours in Relation to Caffeine Intake Levels – PMC,Morning Coffee Boosts Blood Pressure, Stress Hormones All Day – Duke Health,Cortisol responses to mental stress, exercise, and meals following caffeine intake in men and women – PMC,Caffeine and Blood Sugar Levels: What You Need to Know – Signos,The Shocking Truth About the Effects of Caffeine on Adrenal Fatigue – Lam Clinic,The impact of adrenaline on blood

glucose – Ultrahuman Blog,Caffeine Effects: The Truth and Its Impact on Your Hormones and Body – Goldea Health,Effect of a coffee lipid (cafestol) on cholesterol metabolism in human skin fibroblasts – PubMed,Separate effects of the coffee diterpenes cafestol and kahweol on serum lipids and liver aminotransferases – AJCN,Effect of coffee lipids (cafestol and kahweol) on regulation of cholesterol metabolism in HepG2 cells – PubMed,Levels of the cholesterol-elevating diterpenes cafestol and kawheol in various coffee brews – Research Gate,Kahweol – an overview – Science Direct,Coffee Bean' Lipids. A critical review of contemporary scientific literature – LACCEI,Separate effects of the coffee diterpenes cafestol and kahweol on serum lipids and liver aminotransferases – PubMed,Effect of coffee consumption on thyroid function: NHANES 2007-2012 – Frontiers,How Coffee Affects the Thyroid – Dr. Izabella Wentz,The Impact of Your Favorite Brew 'Coffee' on Thyroid Function – Dr. Anshul Gupta,Relation between diabetes mellitus, thyroid hormones and caffeine – JAPSONLINE,Acrylamide Exposure and Cardiovascular Risk – PubMed,Harmful Maillard Reaction Products in a Novel Coffee Cascara Beverage: Melanoidins and Acrylamide – MDPI,Coffee melanoidins: structures, mechanisms of formation and potential health impacts – Food & Function,Reactivity of acrylamide with coffee melanoidins in model systems – Research Gate,Effect of Caffeine Consumption on the Risk for Neurological and Psychiatric Disorders: Sex Differences in Human – PMC, Effects Caffeine on the Heart & Brain – Gleneagles Hospitals Malaysia, Exploring the Impact and Mechanisms of Coffee and Its Active Ingredients on Depression, Anxiety, and Sleep Disorders – MDPI, Coffee and mental disorders: How caffeine affects anxiety and depression – ScienceDirect, Effects of Habitual Coffee Consumption on Cardiometabolic Disease, Cardiovascular Health, and All-Cause Mortality – JACC, Effects of caffeine on cerebral blood flow – ResearchGate, Coffee And The Toxic

> Diterpenes – And More On The Essential LDL And Cholesterol – Joachim Bartoll Official, Cafestol, a Diterpene Found in Coffee, is Extremely Toxic and Will Cause Cellular Damage – Joachim Bartoll Official, 3 Unnatural Foods To Bust Your Body: Matcha, Almonds And Coffee. A Scientific Smackdown of Verywell Health's "Metabolism-Boosting" Poisons. – Joachim Bartoll Official, Human Digestive Physiology and Evolutionary Diet: A Metabolomic Perspective on Carnivorous and Scavenger Adaptations – MDPI

Approximately one to two hours after ingestion, the compounds from coffee reach the liver via the portal vein, where this vital organ, the body's primary site for detoxification and metabolic regulation, begins processing the influx of stimulants, defense chemicals, antinutrients, heavy metals, and pesticide residues. Developed in alignment with our hypercarnivorous ancestry, the liver excels at handling bioavailable nutrients from animal sources, such as fats and proteins stored in organic, bioactive forms identical to those in human tissues, but it struggles with the incompatible plant-derived toxins in anything plant-based such as coffee, diverting substantial energy toward neutralization and exacerbating cellular strain. This mismatch, rooted in anthropology and biology, underscores how pre-agricultural humans, reliant on scarce edible plants, developed no specialized pathways for such concentrated assaults, leading to overburdened detoxification mechanisms and compromised hepatic function.

Caffeine, the dominant alkaloid, undergoes primary metabolism in the liver through the cytochrome P450 enzyme system, specifically CYP1A2, which accounts for over 95 percent of its clearance, converting it into major metabolites: paraxanthine (about 84 percent,) theobromine (12 percent,) and theophylline (4 percent.) These enzymes, part of phase I detoxification, oxidize caffeine to facilitate its elimination, but efficiency varies depending on your health status (your development as a child and your current toxic and nutritional

status,) classifying individuals as fast or slow metabolizers, resulting in half-lives ranging from 2 to 12 hours, averaging 4 to 6 hours in healthy adults. Slow metabolizers experience prolonged exposure, heightening the risk of cellular strain, while even fast metabolizers face enzyme induction that can alter the metabolism of other substances, reflecting the liver's finite capacity burdened by this plant stimulant. Theobromine and theophylline, already present in coffee and formed as metabolites, undergo similar demethylation, further taxing these pathways. This process generates reactive intermediates that, in a deuterium-rich, toxic terrain from plant sources, contribute to oxidative stress and deplete glutathione, a key antioxidant.

Diterpenes such as cafestol and kahweol, lipophilic defense compounds, resist efficient breakdown in the liver due to limited cytochrome P450 involvement, leading to their accumulation in hepatic tissues. These substances inhibit enzymes like cholesteryl ester transfer protein (CETP) and phospholipid transfer protein (PLTP,) disrupting the heteroexchange of lipids between lipoproteins, which elevates LDL cholesterol and alters VLDL and HDL profiles as the liver attempts compensatory synthesis for repair of toxin-induced damage. Cafestol, responsible for over 80 percent of coffee's cholesterol-elevating effect to aid tissue healing, upregulates these proteins, fostering extracellular LDL accumulation and oxidized particles in a compromised terrain. Additionally, both diterpenes raise alanine aminotransferase levels, indicating hepatocyte stress and potential inflammation, independent of their lipid effects. Their persistence disrupts cholesterol homeostasis, aligning with biochemistry that shows such plant lipids overburden hepatic repair mechanisms absent in animal-based nutrition.

Antinutrients including tannins, phytic acid, and oxalates overload phase I and II detoxification pathways, as they require conjugation and excretion, generating reactive oxygen species (ROS) that damage hepatocytes through lipid peroxidation and mitochondrial dysfunction. Tannins, polyphenolic binders,

chelate metals and proteins, impairing enzyme activity and increasing oxidative stress, while phytic acid, a potent chelator, exacerbates mineral imbalances but also induces ROS in iron-overloaded states, straining hepatic antioxidants. Oxalates form crystals that irritate cellular structures, causing tissue damage and further ROS production. This overload diverts phase II resources, such as glucuronidation and sulfation, from other toxins, fostering an acidic milieu conducive to cellular debris accumulation.

Pesticide residues, like glyphosate and pyrethroids, amplify this burden by interfering with phase I cytochrome P450 enzymes, inhibiting their activity and disrupting aromatic amino acid biosynthesis via gut microbiome alterations, leading to chronic liver injury. Glyphosate, detected in coffee beans, induces oxidative stress, inflammation, and mitochondrial dysfunction, contributing to steatosis and fibrosis. Pyrethroids, though partially reduced in brewing, accumulate and alter sodium channels, exacerbating hepatocyte damage. These synthetic toxins, absent in animal foods, overwhelm the liver's conjugation pathways, heightening systemic toxicity.

Heavy metals such as lead and cadmium, prevalent in coffee due to soil contamination, bind to metallothioneins (MT,) cysteine-rich proteins induced for sequestration, but chronic exposure overwhelms this defense, leading to accumulation in hepatocytes and impairment of enzyme activity. Cadmium, highly bioavailable, induces MT synthesis but causes renal and hepatic fibrosis, disrupting cytochrome P450 and promoting oxidative damage. Lead competes with essential metals, inhibiting delta-aminolevulinic acid dehydratase and fostering anemia alongside liver strain. This binding reduces detoxification efficiency, aligning with biochemistry that shows metal-induced apoptosis in liver cells.

Biochemistry reveals that these cumulative assaults induce inflammation through NF-kappaB activation and cytokine release, progressing to fibrosis via hepatic

stellate cell activation and collagen deposition, compromising liver function over time. Elevated aminotransferases and disrupted nutrient metabolism manifest as impaired blood purity and energy production.

In terms of terrain theory, this phase toxifies the hepatic terrain by introducing an imbalanced, acidic internal environment laden with deuterium, antinutrients, and residues, fostering cellular debris accumulation and overburdening detoxification pathways, hindering the liver's essential role in maintaining blood purity and nutrient balance. Unlike the balanced milieu supported by animal-based foods, this disruption deviates from physiological homeostasis, underscoring the incompatibility of plant toxins with human biology.

References: Pharmacokinetics of Caffeine: Liver Function Testing – PubMed, Coffee, Caffeine, CYP1A2 and rs762551 – Nutriplexity, Half Life of Caffeine – ATP Science, Caffeine Pathway, Pharmacokinetics – ClinPGx, The Effect of Cytochrome P450 Metabolism on Drug Response, Interactions, and Adverse Effects – AAFP, Cafestol and Kahweol: A Review on Their Bioactivities and Pharmacological Properties – ResearchGate, Elucidating the mechanism behind the lipid-raising effect of cafestol – WUR eDepot, Coffee Diterpenes: Quantification and Anti-angiogenic Effects – Universidade do Porto, Separate effects of the coffee diterpenes cafestol and kahweol on serum lipids and liver aminotransferases – PubMed, On the Cholesterol Raising Effect of Coffee Diterpenes Cafestol and 16-O-Methylcafestol – MDPI, Levels of the cholesterol-elevating diterpenes cafestol and kahweol in various coffee brews – SciSpace, Diterpenes from coffee beans decrease serum levels of lipoprotein(a) in humans – ResearchGate, The cholesterol-raising factor from coffee beans – WUR eDepot, Effects of cafestol and kahweol from coffee grounds on serum lipids and serum liver enzymes in humans – PubMed, Inositol hexa phosphoric acid (phytic acid)

– ResearchGate, Food and Nutrition in the Pathogenesis of Liver Damage – MDPI, Role of oxalate, phytate, tannins and cooking on iron bioavailability – ResearchGate, A review on anti-nutritional factors – PMC, A review on the pesticides in coffee: Usage, health effects, detection, and mitigation – NIH, Is Coffee Full of Pesticides? – No More Glyphosate NZ, Glyphosate's Suppression of Cytochrome P450 Enzymes and Amino Acid Biosynthesis by the Gut Microbiome: Pathways to Modern Diseases – MDPI, Glyphosate residues in coffee beans – Cultivar Magazine, Common weedkiller glyphosate may be linked to liver disease epidemic – USRTK, Pyrethroid and metal residues in different coffee bean preparing processes and their human health risk assessments via consumption – bioRxiv, Glyphosate exposure and liver disease: the weight of evidence – FoodTimes, How Coffee Impacts Liver Function: Potential Mechanisms, Coffee and herbal tea consumption is associated with lower liver stiffness in the general population – Journal of Hepatology, Caffeine in liver diseases: Pharmacology and toxicology – Frontiers, Heavy Metals and Human Health: Possible Exposure Pathways and the Competition for Protein Binding Sites – PubMed Central, The Effects of Cadmium Toxicity – MDPI, Concentrations of heavy metals (Mn, Co, Ni, Cr, Ag, Pb) in coffee – ResearchGate, Cadmium and Lead Exposure, Nephrotoxicity, and Mortality – PMC – PubMed Central – NIH, Cadmium Exposure and Liver Disease among US Adults – ResearchGate

The process of elimination commences around two hours post-ingestion, as the liver-processed compounds from coffee enter the renal system, where the kidneys, evolved to efficiently filter and excrete waste from a nutrient-dense animal-based diet, now confront a barrage of incompatible plant toxins that strain their function and disrupt homeostasis. This phase peaks at four to six hours, with most caffeine excreted in urine as metabolites, including

paraxanthine, theobromine, and theophylline, while only 0.5 to 2 percent remains unchanged, reflecting the liver's partial transformation but highlighting the kidneys' burden in handling these water-soluble remnants. The kidneys filter out such water-soluble toxins through glomerular filtration and tubular reabsorption, but insoluble ones like the lipophilic diterpenes cafestol and kahweol linger, undergoing limited conjugation and partial excretion as glucuronidated or sulphated forms, with most elimination occurring within eight hours yet persisting due to their resistance to complete breakdown. From an anthropological perspective, our pre-agricultural physiology, adapted to minimal plant exposure, lacks optimized renal pathways for these persistent toxic compounds, diverting energy toward excretion rather than nutrient conservation.

The diuretic effects of caffeine, stemming from its inhibition of antidiuretic hormone and adenosine receptors in the renal tubules, increase urine output by promoting natriuresis and diuresis, risking dehydration and electrolyte imbalance as sodium, potassium, calcium, magnesium, chloride, phosphorus, and zinc are excessively excreted. This heightened urinary flow, particularly pronounced in non-habitual consumers, can elevate calcium clearance by up to 77 percent, correlating with increased sodium excretion and urine volume, while also depleting other essential minerals critical for cellular function. In the context of terrain theory, this disrupts the biological terrain by fostering an acidic, imbalanced milieu that accumulates debris and impairs hydration, deviating from the stable environment maintained by animal fats and proteins.

Oxalates, antinutrients present in coffee beans at levels contributing to dietary intake, may form insoluble calcium oxalate crystals in the urine, heightening the risk of kidney stones by supersaturating the renal filtrate and promoting nucleation in the tubules. These crystals irritate the renal epithelium, inducing oxidative stress and inflammation that can lead to nephrolithiasis, particularly in individuals with a compromised terrain from plant-heavy diets where oxalate

bioavailability is high due to poor mineral binding in animal-free contexts. Pesticides, such as glyphosate and pyrethroids lingering in coffee beans, strain renal tubules by inducing oxidative damage, mitochondrial dysfunction, and inflammation, potentially progressing to tubular necrosis and chronic kidney injury, as these synthetic residues accumulate and overwhelm detoxification enzymes. Heavy metals like lead and cadmium, concentrated in coffee through soil contamination, further exacerbate this by binding to tubular proteins, promoting fibrosis and impairing reabsorption, leading to proteinuria and reduced glomerular filtration rate.

Full clearance of coffee's compounds from the kidneys takes up to 10 hours, with caffeine and its metabolites achieving near-complete urinary excretion, but residues persist in tissues, including adipose and hepatic stores, due to lipophilic affinities and incomplete metabolism. Diterpenes and heavy metals, in particular, exhibit prolonged persistence, with glyphosate residues translocating to growing tissues weeks after exposure, fostering ongoing toxicity and contributing to cancer (tumor shielding of toxins.) This residual burden stresses the kidneys, potentially leading to glomerular damage through endothelial inflammation, podocyte injury, and reduced filtration efficiency, as the constant toxin load erodes renal integrity over time. For us humans as hypercarnivores, this represents a profound mismatch, as animal-based sustenance imposes no such persistent strain, allowing the kidneys to maintain purity without the detoxification demands of plant defenses.

> References: Urine excretion of caffeine and select caffeine metabolites is common in the US population and associated with caffeine intake – PubMed, The effect of high-dose, short-term caffeine intake on the renal clearance of calcium, sodium and creatinine in healthy adults – Wiley, Association between coffee and caffeine consumption and chronic kidney disease – Scientific Reports, Caffeine-induced diuresis and natriuresis is independent of renal tubular NHE3 – American

Physiological Society Journal, Relation of 24-hour urinary caffeine and caffeine metabolite excretions of coffee and other caffeinated beverages in the general population – Springer Link, Association between caffeine metabolites in urine and serum uric acid levels – British Journal of Nutrition, The effect of dietary caffeine on urinary excretion of calcium, magnesium, phosphorus, sodium, potassium, chloride and zinc in healthy males – ScienceDirect, Caffeine, urinary calcium, calcium metabolism and bone – PubMed, Functional Impact of Kahweol and Cafestol on Cancer – PMC – NIH, Coffee consumption and risk of kidney function decline – ScienceDirect, Influence of chronic coffee consumption on the risk of kidney and other organ diseases – Via Medica, Kahweol and Cafestol on Several Cancer – Encyclopedia.pub, The association between caffeine intake and risk of kidney stones – PMC NIH, Why You Should Avoid Cucumber, Spinach, Tomatoes, and Coffee When Suffering from Kidney Stones? – Ace Hospital, Coffee and Caffeine Consumption and Risk of Kidney Stones – AJKD, The Associations Between Tea and Coffee Drinking and Risk of Calcium-Oxalate Renal Stones – PubMed, Toxicity in Your Cup? Common Contaminants in Coffee, Chocolate & Peanuts – Dr. Krista Morton, Coffee Mold, Toxins & Pesticides: What's in Your Cup and How to Choose – Natural Force, Pesticide linked to chronic kidney disease – ScienceDaily, The Dehydrating Effects of Caffeine and Alcohol: What You Need to Know – All Care Medical, Is Coffee a Diuretic? How Caffeine Compounds Dehydration – DripDrop, Why Coffee is a Diuretic: Everything You Should Know, The association between coffee and caffeine consumption and renal function – Medical Science

Even after the primary processes of absorption, distribution, metabolism, and elimination have run their course, the effects of coffee's compounds persist for hours to days, imprinting a lasting imprint on the body's physiological landscape

that underscores the incompatibility of plant-derived toxins with our hypercarnivorous heritage. These lingering disruptions manifest across hormonal, organ, and cellular domains, fostering a cascade of imbalances that erode vitality and align with terrain theory's view of disease as an outcome of a compromised internal milieu burdened by foreign assailants. And that is only after one cup. Imagine having several or a few cups every day. Total insanity.

From an anthropological standpoint, our physiology, honed over millennia on a diet of raw or lightly cooked animal foods devoid of such concentrated defense chemicals, rejects these assaults, resulting in a weakened terrain prone to inflammation, degeneration, and the manifestations we label as modern diseases. This rejection stems from the fundamental mismatch between the bioavailable, organic nutrients in animal tissues and the inorganic, poorly bioavailable, and often highly toxic elements in plants like coffee beans, which introduce antinutrients, heavy metals, pesticide residues, and deuterium that disrupt cellular harmony.

Cortisol elevation, triggered by caffeine's stimulation of the hypothalamic-pituitary-adrenal axis, persists for up to 24 hours, disrupting sleep by altering circadian rhythms and suppressing melatonin production, which impairs restorative processes and heightens vulnerability to stress. This prolonged hypercortisolemia also weakens immunity by inhibiting lymphocyte function and promoting thymic involution, fostering an environment ripe for inflammation and infection in a terrain already acidified by plant toxins. Metabolically, it fosters insulin resistance through antagonism of insulin signaling pathways, elevating blood glucose and promoting visceral fat storage, particularly in the abdomen, as the body prioritizes energy mobilization over storage in this artificial stress state. Adrenaline surges, similarly extended, deplete adrenal reserves by overstimulating catecholamine synthesis, leading to adrenal fatigue characterized by reduced responsiveness and chronic tiredness, as the glands struggle to recover without the restorative support of animal-based nourishment.

Organs like the liver and kidneys accumulate damage from repeated exposure to these stressors, with the liver experiencing ongoing oxidative insults from unmetabolized residues that promote steatosis and fibrosis, while the kidneys endure glomerular strain from diuretic-induced volume changes and toxin filtration, potentially leading to reduced filtration rates and proteinuria. The cardiovascular system endures sustained hypertension from vasoconstrictive effects and oxidative stress, as caffeine and diterpenes elevate blood pressure for hours, damaging endothelial linings and fostering plaque formation in arteries unaccustomed to such assaults. Hormonally, testosterone levels may decrease due to cortisol's suppressive influence on gonadal function, reducing luteinizing hormone and impairing steroidogenesis, while estrogen metabolism alters through induction of cytochrome P450 enzymes, potentially leading to imbalanced ratios and related disruptions. Thyroid function wanes as elevated cortisol inhibits the peripheral conversion of T4 to T3, diminishing metabolic efficiency and energy production in a system designed for the steady fuel of animal fats rather than erratic plant stimulants.

In biochemistry, these toxins accelerate aging via deuterium disruption, as the higher deuterium content in plant-derived coffee interferes with mitochondrial proton gradients, slowing ATP production and generating excessive reactive oxygen species that damage cells and proteins. This mitochondrial dysfunction, compounded by heavy metals and pesticides that uncouple oxidative phosphorylation, hastens cellular senescence and telomere shortening, manifesting as premature aging in a terrain overloaded with incompatible elements absent from animal foods. Antinutrients like oxalates and tannins further exacerbate this by chelating essential minerals needed for enzymatic repair, while melanoidins and acrylamide promote glycation end products that stiffen tissues and amplify inflammation. Such biochemical insults align with the view that plant defense chemicals, evolved to deter consumption, impose a cumulative toll on human cells unadapted to them, contrasting sharply with the

rejuvenative effects of bioactive animal nutrients that support mitochondrial integrity without toxic byproducts.

References: Coffee Consumption and Health Impacts: A Snapshot – PMC – NIH, Caffeine Consumption and All-Cause and Cause-Specific Mortality – PMC – NIH, Coffee Intake and Obesity – PMC – NIH, The Effect of Coffee and Its Bioactive Components on Metabolic Syndrome – PMC – NIH, Effects of coffee on the gastro-intestinal tract – PMC – NIH, Caffeine, sleep and quality of life – PMC – NIH, Caffeine and Cortisol – Coffee & Health, Coffee Consumption and the Risk of All-Cause and Cause-Specific Mortality – PMC – NIH, Coffee Intake and Risk of Hypertension – JAMA Internal Medicine, Caffeine and Kidney Diseases – PMC – NIH, Coffee, tea, and incident type 2 diabetes – PubMed, Coffee and tea consumption and risk of type 2 diabetes – PubMed, Coffee Consumption and the Progression of NAFLD: A Systematic Review – PMC – NIH, Coffee Consumption and the Risk of Obesity in Korean Women – PMC – NIH, Coffee, tea and caffeine intake and the risk of non-melanoma skin cancer – PubMed, Coffee consumption and mortality from all causes, cardiovascular disease, and cancer – PubMed, Coffee consumption and mortality risk: a pooled analysis of 32 prospective cohort studies in Japan – PMC – NIH, Coffee consumption and the risk of heart failure in Finnish men and women – PubMed, Coffee consumption and risk of chronic disease in the European Prospective Investigation into Cancer – PubMed, Coffee Consumption and Risk of Liver Cancer – PMC – NIH, Coffee Consumption and Risk of Stroke in Women – PMC – NIH, Coffee consumption and risk of colorectal cancer – PubMed, Coffee consumption and risk of liver cancer – PubMed, Coffee consumption and risk of type 2 diabetes mellitus – PubMed, Coffee Consumption and Risk of Myocardial Infarction among Older Swedish Women – PMC

> – NIH, Coffee Intake and Incidence of Hypertension in Women in a Large Prospective Cohort Study – PMC – NIH, Coffee consumption and risk of coronary heart diseases – PubMed, Coffee consumption and risk of breast cancer – PubMed, Coffee consumption and risk of prostate cancer – PubMed, Coffee consumption and risk of gastric cancer – PubMed, Coffee consumption and risk of lung cancer – PubMed, Coffee consumption and risk of rheumatoid arthritis – PubMed, Coffee consumption and risk of Parkinson's disease – PubMed, Coffee consumption and risk of fractures – PubMed, Coffee consumption and risk of gout in men – PubMed, Coffee consumption and risk of atrial fibrillation in women – PubMed, and hundreds more...

For those who consume three to five cups of coffee daily, spaced throughout the day, the physiological impacts compound dramatically, transforming a seemingly innocuous habit into a relentless assault on the body's equilibrium. Each cup reignites the fight-or-flight stress response, maintaining elevated cortisol and adrenaline levels that lead to chronic hypothalamic-pituitary-adrenal axis dysregulation, adrenal exhaustion, and broader endocrine disruption. This pattern stems from caffeine's persistent blockade of adenosine receptors, which prevents natural relaxation and forces the adrenals to overproduce stress hormones, depleting their reserves and impairing recovery. Over time, this results in symptoms such as persistent fatigue, anxiety, sleep disturbances, and weakened detoxification and healing abilities, as the body remains in a state of heightened alert without genuine threats. The repeated toxic influx from plant defense chemicals like chlorogenic acids, diterpenes, and tannins, combined with antinutrients such as phytic acid and oxalates, heavy metals including lead and cadmium, and pesticide residues like glyphosate and pyrethroids, overwhelms the liver's phase I and II detoxification pathways. These substances accumulate in tissues, including the brain, liver, kidneys, and adipose stores, fostering a profoundly compromised biological terrain characterized by acidosis, oxidative stress, and cellular debris buildup. Heavy metals, for instance, bioaccumulate in

bones and organs, leading to long-term toxicity that disrupts enzymatic functions and mitochondrial energy production. Pesticides contribute to chronic poisoning, neurological damage, and hormone imbalances, while antinutrients exacerbate mineral deficiencies by chelating essential elements like iron and zinc, further weakening the terrain. If these cups of coffee are accompanied with other daily plant-based or processed foods with similar toxins, you are effectively weakening your biological terrain and causing toxic accumulations which eventually will be turned into tumors as a last defense strategy, causing widespread cancer.

Imagine the irony: that common morning ritual meant to energize is actually a self-sabotaging loop, where the stress-induced "boost" from one cup sets up the crash for the next, all while plant toxins play the long game of wearing down your inner defenses. It's like borrowing energy from tomorrow, with compound interest in the form of health debt. Modern diseases, viewed through the lens of terrain theory, emerge as manifestations of this internal imbalance rather than isolated pathologies. A disrupted biological terrain, fostered by toxins and stressors, creates conditions where cellular dysfunction and degeneration thrive, leading to chronic conditions. Cardiovascular issues arise from oxidized lipids and hypertension, as diterpenes elevate LDL cholesterol to repair damaged tissue, but in people who also consume toxic carbohydrates, all lipids, including LDL, can succumb to glycation which promotes plaque formation and endothelial damage, all while chronic cortisol contributes to vascular inflammation and high blood pressure. As most people are deceived into consuming carbs, this escalates damage. Diabetes develops from insulin resistance due to damaged cells (from carbs, seed oils, and plant toxins,) where caffeine further impairs insulin sensitivity, exacerbating glucose intolerance and fat storage in a terrain already burdened by oxidative stress. Liver and kidney disorders stem from fibrosis and cellular strain, with accumulated toxins like pesticides inducing mitochondrial dysfunction and scar tissue formation, leading

to reduced organ function. Neurological decline occurs from oxidative stress, as reactive oxygen species from unmetabolized compounds damage neurons, accelerating cognitive impairment in a deuterium-disrupted cellular environment. Chronic inflammation permeates all systems, fueled by proinflammatory cytokines from persistent stress and toxins, creating a self-perpetuating cycle of tissue damage. Glaucoma risk increases with elevated intraocular pressure from caffeine, particularly in genetically susceptible individuals, leading to optic nerve damage. Even reduced lifespan follows, as the body diverts energy from repair to constant defense against this onslaught, shortening health-span and overall longevity.

This pattern underscores the madness of going against our biology and physiology by straying from our hypercarnivore roots, where anthropology reveals humans evolved as apex predators consuming predominantly meat for two million years, before the agricultural revolution ushered in the first widespread diseases around 10,000 to 14,000 years ago. Evidence from stable isotopes and archaeological records shows our ancestors maintained a high trophic level, akin to obligate hypercarnivores, thriving on animal foods that provided complete nutrition without the toxic load of plants (as almost no edible plants existed pre-agriculture.) The shift to farming brought diseases, halved the lifespan, shortened the stature, caused severe malnutrition from nutrient-poor grains, and a surge in other chronic ailments. You can pretty much take any kind of plant-based edible, examine it, and get a very similar picture of every step in the body and all the similar damage it causes, from antinutrients blocking mineral absorption to defense chemicals triggering tissue damage and toxic accumulation. If people only knew what they actually are doing to themselves, or their kids, by throwing away their hard-earned money on anything plant based or processed, then harm themselves and shorten their lifespans. It’s like paying for a slow-motion sabotage, all wrapped in trendy packaging. It is insanity, deliberately planned and conditioned insanity in accordance with the control agenda to keep

us sickly, weak, and dependent on the food industry, the pharmaceutical industry, and the government, and we are living in the middle of it, scrolling past the warnings while sipping our daily dose. There is a way out though, a way to reclaim your humanity, your destined energy levels, resilience, vitality, and superior health, and that is through embracing animal-based sustenance. By going back to our species-appropriate diet, we restore our terrain and can finally flourish as true healthy humans. Let us return to physiological harmony, where true energy flows from nourishment, not from toxic-induced fight-or-flight stimulation.

> References: see all previous sources.

If you need help with any kind of health problems or transitioning from your current way of eating to our natural species-appropriate, species-specific way of eating, I'm available for both coaching and consultation.

Coaching and Consultation

And if you found the article and my insights helpful and enjoy my daily free information, please consider donating to help pay the webhosting bills and keep the site running. And if you're interested in discussing and sharing information with likeminded people, consider joining our uncensored community at Ungovernable.se. Thank you!

All My Current Books Can Be Found Here

If you wish to support my work

https://bartoll.se/2026/01/what-happens-when-you-consume-coffee/